



Dermal Filler Pre & Post Care

Pre-Treatment Instructions

- Schedule your injection at a time when minor swelling or bruising will not disrupt your social obligations. Each time you are treated, outcomes may vary.
- To lessen the likelihood of bleeding or bruising, discontinue use of blood thinning products for 10 days prior to treatment, if approved by your primary care provider. This includes: aspirin, ibuprofen (Motrin or Advil), naproxen, Aleve, fish oil, St. John's Wort, flax, Vitamin E, garlic, Ginkgo Biloba, ginseng, etc. Alcohol is also to be avoided 24 hours pre- or post- injectable treatment, as it is also a blood thinner.
- Please notify your provider if you have any history of cold sores, as we will consider pre-medicating with an antiviral prescription prior to injections. Cancel your appointment if you have any open sores in the treatment area.
- Any injection introduces the risk for infection. For this reason, the following is taken into consideration when proceeding with injections. We will delay injectables if you:
 - Have had any dental procedures in the past 30 days or are anticipating dental procedures or cleanings in the next 30 days.
 - Have had a surgical procedure in the past 30 days or are anticipating a surgery or procedure.
 - Have had any immunizations in the past 30 days or if you're anticipating immunizations.
 - Have had any tattooing or permanent makeup within the treatment area within the last 30 days.
 - Take any immunosuppressive/injectable medications (ex: Biologics such as Humira, Skyrizi, Stelara, Enbrel, Taltz, Cosentyx, Tremfya).
 - Additional contraindications: Pregnancy, breastfeeding, allergies to components of dermal fillers, open sores in areas to be treated.

Post-Treatment Instructions

- Swelling, bruising, tenderness, numbness, and areas of firm nodules may occur.
- Cosmetic fillers are long lasting, but not permanent. Longevity depends on the areas treated and your body's metabolism.
- DO NOT APPLY ANYTHING TO THE SKIN UNTIL THE DAY AFTER TREATMENT: No cleanser, moisturizer, or makeup!
- DO NOT massage, touch, or manipulate the injection site.
- Ice packs may be used to the treated area during the first 12 hours. Ice for 15 minutes on every hour.
- Avoid dental work for 30 days after filler injections.
- Avoid alcohol and heating your core body temp for 24-48 hrs. (intense work outs, sauna, hot tub, etc.)
- If experiencing swelling, sleep with head elevated for 2-3 days to decrease swelling.
- You may take acetaminophen/Tylenol if you experience any mild tenderness or discomfort.
- Wait a minimum of 2 weeks (or as directed by your provider) before receiving any skin care or laser treatments.
- *If you experience any of the following symptoms, contact our office immediately: dusky or white discoloration of injected areas, mottling or unusual bruising, severe or increasing pain, redness, increasing warmth or coolness to touch, severe swelling, itching, blisters, difficulty swallowing or smiling, fever, or chills. If you notice onset of any of these symptoms, call our office immediately at (530)559-8558 to speak with a provider.*